arXiv:1110.1820v1 [math.DG] 9 Oct 2011

ON THE GEOMETRY OF FOUR DIMENSIONAL RIEMANNIAN MANIFOLD WITH A CIRCULANT METRIC AND A CIRCULANT AFFINOR STRUCTURE

DIMITAR RAZPOPOV

Abstract. We consider a 4-dimensional Riemannian manifold M with a metric g and an affinor structure q. We note the local coordinates of g and q are circulant matrices. Their first orders are (A, B, C, B), $A, B, C \in FM$ and $(0, 1, 0, 0)$, respectively.

Let ∇ be the connection of g. Further, let $\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6$ be the sectional curvatures of 2-sections $\{x, qx\}$, $\{x, q^2x\}$, $\{q^3x, x\}$, $\{qx, q^2x\}$, $\{qx, q^3x\}$, $\{q^2x, q^3x\}$ for arbitrary vector $x \in T_pM$, $p \in M$. Then we have that $q^4 = E$; $\quad g(qx, qy) = g(x, y), \quad x,\ y \in \chi M$.

The main results of the present paper are

1) There exist a q-base in T_pM, $p \in M$.

2) if $\nabla q = 0$, then $\mu_1 = \mu_3 = \mu_4 = \mu_6$, $\mu_2 = \mu_5 = 0$,

Mathematics Subject Classification (2010): 53C15, 53B20

Keywords: Riemannian metric, affinor structure, sectional curvatures

1. Introduction

The main purpose of the present paper is to continue the considerations on the differential geometry of Riemannian manifolds with a circulant metric g, and a circulant affinor structure q [1], [2], [3], [4].

Let M be a Riemannian manifold with a metric g and affinor structure q, such that [4]:

$$g_{ij} = \begin{pmatrix} A & B & C & B \\ B & A & B & C \\ C & B & A & B \\ B & C & B & A \end{pmatrix}, \quad q_i^{\cdot j} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \tag{1}$$

where $det g_{ij} = (A - C)^2((A + C)^2 - 4B^2)$, and $A = A(p)$, $B = B(p)$, $C = C(p)$, $p = (x^1, x^2, x^3, x^4) \in D \subset R^4$ are smooth functions in the local coordinate system $\{x^1, x^2, x^3, x^4\}$. We suppose

$$0 < B < C < A. \tag{2}$$

By virtue of [2] we easily get that g is positively defined metric.

We know from [4] that

$$q^4 = id, \qquad q \neq \pm id, \quad q^2 \neq \pm id \tag{3}$$

$$g(qx, qy) = g(x, y), \qquad x,\ y \in \chi M. \tag{4}$$

Let ∇ be the Levi-Chivita connection of g and R be the curvature of ∇, i.e $R(x,y)z = \nabla_x \nabla_y z - \nabla_{[x,y]} z$. We consider the associated with R tensor field R of type $(0,4)$, defined by the condition

$$R(x,y,z,u) = g(R(x,y)z,u), \qquad x,y,z,u \in \chi M.$$

Then we have

$$\nabla q = 0 \Leftrightarrow \mathrm{grad} A = \mathrm{grad} C.q^2, \mathrm{grad} B = \mathrm{grad} C.(q+q^3) \tag{5}$$

$$\nabla q = 0 \Rightarrow R(x,y,qz,qu) = R(x,y,z,u), x,y,z,u \in \chi M. \tag{6}$$

Further in our considerations we suppose that M owns all properties (1)–(6).

2. q-BASES

Theorem 2.1. *Let M be a 4-dimensional Riemannian manifold with a metric g and an affinor structure q with local coordinates (1). If p is a point in M and x is an arbitrary vector in T_pM then there exist a q-base in T_pM.*

Proof. Let $p \in M$, $x = (x^1, x^2, x^3, x^4) \in T_pM$, and $qx \neq x$, $q^2x \neq x$. Then using (1) we easily see that the determinant of the coordinates of x, qx, q^2x and q^3x is not zero. So, $\{x, qx, q^2x, q^3x\}$ is a q-base in T_pM. □

Theorem 2.2. *Let M be a 4-dimensional Riemannian manifold with a metric g and an affinor structure q with local coordinates (1). If p is a point in M and x is an arbitrary vector in T_pM then there exist an orthonormal q-base in T_pM.*

Proof. Let $p \in M$, $x = (x^1, x^2, x^3, x^4) \in T_pM$. We suppose that

$$x^4 = 0, \qquad x^2 = \frac{\sqrt{A+B-2C} - \sqrt{A+B+2C}}{2\sqrt{A+B-2C}\sqrt{A+B+2C}},$$

and x^1 and x^3 satisfy the following system

$$\begin{aligned} x^1 + x^3 &= \frac{\sqrt{A+B-2C} - \sqrt{A+B+2C}}{2\sqrt{A+B-2C}\sqrt{A+B+2C}}, \\ x^1 x^3 &= \frac{2B^2 - C^2 - AC}{2(A-C)\sqrt{A+B-2C}\sqrt{A+B+2C}}. \end{aligned} \tag{7}$$

The solutions of the system (7) are solutions of the square equations with discriminant

$$D = (x^1+x^3)^2 - 4x^1x^3 = \frac{A^2 + C^2 + 2AC + (A-C)\sqrt{(A+C)^2 - 4B^2} - 4B^2}{2A - C((A+C)^2 - C) - 4B^2}.$$

We have $D > 0$ because of (2).

Now we can verify directly that $g(x,x) = 1$, $g(x,qx) = 0$, $g(x,q^2x) = 0$ and consequently, on due to (1), we see that $g(x,q^3x) = g(qx,q^2x) = g(qx,q^3x) = g(q^2x,q^3x) = 0$. So, $\{x, qx, q^2x, q^3x\}$ is an orthonormal q-base. □

An example. Let $p \in M$ and $\{x, qx, q^2x, q^3x\}$ be a q-base in T_pM. Let L, N, S, T be four points in T_pM such that $PL = x$, $PN = qx$, $PS = q^2x$, $PT = q^3x$. Let α and β be the corresponding angels between x and qx, and x and q^2x, respectively. Using (1)–(4) we get $\|LN\| = \|LT\| = \|ST\| = 2\|x\|^2(1 - \cos\alpha)$,

$\|LS\| = \|NT\| = 2\|x\|^2(1-\cos\beta)$. We see that $LNST$ is a pyramid with six sides which are all congruent isoseles triangles. The angles of such a side of the pyramid we note by γ and δ, so $2\gamma+\delta=\pi$. They satisfy

$$\cos\gamma = \frac{1-\cos\beta}{2\sqrt{1-\cos\alpha}\sqrt{1-\cos\beta}}, \quad cos\delta = \frac{1-2\cos\alpha+\cos\beta}{2(1-cos\alpha)}.$$

Evidently, in the case when $\{x, qx, q^2x, q^3x\}$ is an orthonormal q-base then the every side of the pyramid is an equilateral triangle.

3. CURVATURES

Let p be a point in M and x, y be two linearly independent vectors on T_pM. It is known the value

$$\mu(E;p) = \frac{R(x,y,x,y)}{g(x,x)g(y,y)-g^2(x,y)} \tag{8}$$

is the sectional curvature of 2-section $E=\{x,y\}$.

For a vector $x=(x^1,x^2,x^3,x^4)$ from T_pM we suppose $((x^1-x^3)^2+(x^2-x^4)^2)(x^1-x^2+x^3-x^4)(x^1+x^2+x^3+x^4)\neq 0$. By using (3) we get that the vectors x, qx, q^2x, q^3x are linearly independent.

Theorem 3.1. *Let M be the Riemannian manifold with a metric g and a parallel structure q, defined by (1) and (2), respectively. Let p be a point in M and x be an arbitrary vector in T_pM. Then the sectional curvatures of 2-sections $E_1=\{x,qx\}$, $E_3=\{q^3x,x\}$, $E_4=\{q^2x,qx\}$, $E_6=\{q^2x,q^3x\}$ are equal among them, and the sectional curvatures of 2-sections $E_2=\{x,q^2x\}$ $E_5=\{q^3x,qx\}$ are vanish.*

Proof. From (6) we find

$$R(x,y,z,u)=R(x,y,qz,qu)=R(x,y,q^2z,q^2u)=R(x,y,q^3z,q^3u). \tag{9}$$

In (9) we get the following substitutions:

a) $z=x$, $y=u=qx$; $x\sim qx$, $z=x$, $y=u=q^2x$; $z=x$, $y=u=q^3x$.

$$R(x,qx,x,qx)=R(x,q^3x,x,q^3x), \qquad R(x,q^2x,x,q^2x)=0. \tag{10}$$

b) $z=x$, $y=qx$, $u=q^2x$; $z=qx$, $y=u=q^2x$; $z=x$, $y=q^3x$, $u=q^2x$; $z=qx$, $y=u=q^3x$.

$$R(x,qx,x,q^2x)=R(x,q^2x,qx,q^2x)=R(x,q^3x,q^2x,x)=R(x,q^3x,qx,q^3x)=0$$

c) $z=x$, $y=qx$, $u=q^3x$; $z=qx$, $y=u=q^2x$; $z=qx$, $y=qx$, $u=q^2x$; $z=q^2x$, $y=qx$, $u=q^3x$.

$$-R(x,qx,x,q^3x)=R(x,qx,qx,q^2x)=R(x,qx,q^2x,q^3x)=R(x,qx,x,qx)$$

d) $z=qx$, $y=qx$, $u=q^3x$; $z=qx$, $y=q^2x$, $u=q^3x$; $z=q^2x$, $y=q^2x$, $u=q^3x$; $z=q^2x$, $y=qx$, $u=q^3x$.

$$R(x,q^2x,qx,q^3x)=R(x,q^2x,q^2x,q^3x)=R(qx,q^2x,qx,q^3x)=0$$

e) $z=qx$, $y=q^3x$, $u=q^2x$; $z=q^2x$, $y=u=q^3x$; $x\sim qx$, $z=q^2x$, $y=q^2x$, $u=q^3x$; $x\sim qx$, $z=q^2x$, $y=u=q^3x$.

$$-R(x,q^3x,qx,q^2x)=-R(x,q^3x,q^2x,q^3x)=R(qx,q^2x,q^2x,q^3x)=R(x,qx,x,qx)$$

and

$$R(qx,q^3x,q^2x,q^3x)=0$$

f) $x \sim qx$, $z = qx$, $y = q^3x$, $u = q^3x$; $x \sim qx$, $z = qx$, $y = q^2x$, $u = q^2x$; $x \sim q^2x$, $z = q^2x$, $y = q^3x$, $u = q^3x$.

$$R(qx, q^3x, qx, q^3x) = 0 \tag{11}$$

$$R(qx, q^2x, qx, q^2x) = R(q^2x, q^3x, q^2x, q^3x) = R(x, qx, x, qx) \tag{12}$$

From (4), (8), (10)–(12) we get

$$\mu(E_1; p) = \mu(E_3; p) = \mu(E_4; p) = \mu(E_6; p) = \frac{R(x, qx, x, qx)}{g^2(x, x) - g^2(x, qx)}.$$

$$\mu(E_2; p) = \mu(E_5; p) = 0.$$

By virtue of the linear independence of x and qx we have

$$g^2(x, x) - g^2(x, qx) = g^2(x, x)(1 - cos\varphi) \neq 0,$$

where φ is the angle between x and qx. □

References

[1] G. Dzhelepov, I. Dokuzova, D. Razpopov. *On a three dimensional Riemannian manifold with an additional structure.* arXiv:math.DG/0905.0801

[2] G. Dzhelepov, D. Razpopov, I. Dokuzova. *Almost conformal transformation on Riemannian manifold with an additional structure*, Proceedings of the Anniversary International Conference - REMIA, Plovdiv, (2010), 125–128, arXiv:math.DG/1010.4975

[3] I. Dokuzova, D. Razpopov. *On affine connections in a Riemannian manifold with a circulant metric and two circulant affinor structure*, Proceedings of the 40-th Anniversary Spring Conference of UBM, Borovets, (2011), 170–176

[4] K. Yano. *Differential geometry.* Pergamont press, New York, 1965

[5] H.Hristov. *Mathematical methods in physics.* Science and Art, Sofia, 1967(in bulgarian)

Dimitar Razpopov
Department of Mathematics and Physics
Agricultural University of Plovdiv
Bulgaria 4000
e-mail:drazpopov@qustyle.bg